



Congenital Cardiology Solutions

SURGICAL MANAGEMENT OF ATRIOVENTRICULAR SEPTAL DEFECTS WITH AORTIC ARCH OBSTRUCTION IN NEONATES: A MULTICENTER STUDY

ACC Moderated Poster Contributions
McCormick Place South, Hall A
Sunday, March 25, 2012, 9:30 a.m.-10:30 a.m.

Session Title: Congenital Cardiology Solutions: Therapy
Abstract Category: 28. Congenital Cardiology Solutions: Therapy
Presentation Number: 1139-303

Authors: John M. Costello, Jeffrey Shuhaiber, Andrew Shin, Jeffrey Gossett, David Wypij, Carl Backer, Frank Hanley, Muhammad S. Khan, Charles Fraser, Frederic Jacques, Glen Van Arsdell, John Mayer, Children's Memorial Hospital, Northwestern University Feinberg School of Medicine, Chicago, IL, USA, Children's Hospital Boston, Harvard Medical School, Boston, MA, USA

Background: For neonates with atrioventricular septal defect (AVSD) and aortic arch obstruction thought to be candidates for a two ventricular repair, surgical options include a primary repair or a staged approach involving initial arch repair followed by delayed AVSD repair. We sought to determine whether a difference in outcomes exists depending on surgical approach.

Methods: In this retrospective cohort study, we included 66 consecutive neonates with AVSD and aortic arch obstruction who underwent surgery prior to 28 days of age at six centers from 1990-2009 (four centers) and 1999-2009 (two centers). Characteristics and outcomes between patients undergoing primary vs. staged repair were compared.

Results: Of 66 study patients, 31 (47%) underwent primary and 35 (53%) underwent staged repair. At initial presentation, compared to the primary repair group, those who underwent staged repair were more likely to receive prostaglandin E1 prior to the first operation (82% vs. 60%, $p=0.047$). There were no other differences in demographic characteristics, cardiac anatomical details, severity of baseline atrioventricular valve regurgitation, or surgical technique. Those undergoing primary repair were more likely to have a least moderate mitral regurgitation early after repair (42% vs. 17%, $p=0.03$), require an unplanned cardiac re-operation during the initial hospitalization or follow-up (45% vs. 15%; $p=0.009$), and tended to have one or more major complications (42% vs. 20%; $p=0.053$). Compared to staged repair patients, those undergoing primary repair tended to be more likely to die during a cardiac surgical hospitalization (26% vs. 9%; $p=0.06$). At a median of 2.0 years (range 0-15.2) of follow-up, overall mortality at follow-up was higher (35% vs. 14%; log rank $p=0.02$). There was no association between later era and surgical approach or mortality.

Conclusions: For neonates with AVSD and aortic arch obstruction, when compared with primary repair, a staged approach involving initial arch repair followed by delayed AVSD repair was associated with lower overall morbidity and mortality.